



Characterization of bifurcating non-linear normal modes in piecewise linear mechanical systems

Paolo Casini, Fabrizio Vestroni*

Dipartimento di Ingegneria Strutturale e Geotecnica (DISG), Università degli Studi di Roma "La Sapienza", Via Eudossiana 18-00184 Roma, Italy

ARTICLE INFO

Article history:

Received 16 October 2009

Received in revised form

6 August 2010

Accepted 17 August 2010

Keywords:

Non-linear normal modes

Piecewise linear systems

Internal resonance

Damaged systems

ABSTRACT

The non-linear modal properties of a vibrating 2-DOF system with non-smooth (piecewise linear) characteristics are investigated; this oscillator can suitably model beams with a breathing crack or systems colliding with an elastic obstacle. The system having two discontinuity boundaries is non-linearizable and exhibits the peculiar feature of a number of non-linear normal modes (NNMs) that are greater than the degrees of freedom. Since the non-linearities are concentrated at the origin, its non-linear frequencies are independent of the energy level and uniquely depend on the damage parameter. An analysis of the NNMs has been performed for a wide range of damage parameter by employing numerical procedures and Poincaré maps. The influence of damage on the non-linear frequencies has been investigated and bifurcations characterized by the onset of superabundant modes in internal resonance, with a significantly different shape than that of modes on fundamental branch, have been revealed.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

The classical modal analysis in linear dynamics can be extended to non-linear systems governed by smooth equations by introducing the concept of non-linear normal modes (NNMs). According to the classical definition given by Rosenberg [1,2], a NNM of an undamped system is defined as a synchronous periodic oscillation where all generalized coordinates of the system reach their extreme values or pass through the zeros simultaneously. The NNMs of a system are important because, in analogy to linear theory, resonance in forced systems typically occurs in the neighborhood of NNM frequencies. Hence, knowledge of the normal modes of a non-linear system can provide valuable insight regarding the position of its resonances, a feature of considerable engineering importance. Moreover, since the number of normal modes of a non-linear system may exceed its degrees of freedom (*superabundant NNMs*), certain forced resonances are essentially non-linear and have no analogies in linear theory; in such cases a linearization of the system might not be possible, or might not provide all the possible resonances that can be experienced.

In recent years, this item has revealed its importance also for piecewise-smooth mechanical systems (PSS): these systems are governed by sets of ordinary differential equations, which are

smooth in regions of phase space; smoothness is lost as trajectories cross the boundaries between adjacent regions (*discontinuity boundaries*). Much effort in science and engineering has focused on these kinds of problems [3]. Typical mechanical applications include: oscillators colliding with a deformable or rigid stop [4,5], block assemblies connected by no-tension springs [6], beams with breathing cracks [7–10] and stick-slip mechanical systems with friction [11,12].

Particular attention has been devoted to the investigation of the non-linear modal properties of these systems. To this end, the classical Rosenberg's definition has been extended, to make it suitable for non-conservative and non-smooth systems: NNM is then defined as any periodic motion in which all generalized coordinates vibrate without necessarily passing through the zeros simultaneously [9,13]. Important contributions to the understanding of NNMs from a general point of view can be found in [8,14–21].

In a recent study [13], a discrete model of a beam with a breathing crack, an example of a non-linearizable system, has been dealt with using the asymptotic method of Lindstedt–Poincaré and limiting the analysis to the fundamental branch solutions and their stability. In [22], a similar system was then numerically investigated by means of the Poincaré map. Particular attention was given to the onset of superabundant modes along the backbones of the two fundamental modes.

The present paper aims to give a general picture of the basic non-linear dynamics of these systems. In particular the onset of superabundant modes at internal resonances will be characterized and it will be shown that the specific features of each

* Corresponding author.

E-mail addresses: p.casini@uniroma1.it (P. Casini), vestroni@uniroma1.it (F. Vestroni).