

Herschel out-of-field stray-light characterization

U. Klaas · K. Okumura · M. Ferlet · T. Müller ·
M. Sanchez-Portal · B. Altieri · D. Doyle ·
G. L. Pilbratt

Received: 1 July 2013 / Accepted: 10 October 2013
© Springer Science+Business Media Dordrecht 2013

Abstract Out-of-field stray-light spots of the *Herschel* telescope optics relative to the PACS and SPIRE instrument apertures were modeled by ray tracing simulations with the Advanced Systems Analysis Program (ASAP, by Breault Research Organization) prior to launch. The predicted stray-light behaviour was verified by dedicated stray-light calibration observations in-flight. This resulted in a special feature of the *Herschel* Science Mission Planning Software, marking the sky positions of stray-light spots by the very bright infrared planetary sources Venus, Mars, Jupiter, and

Herschel is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

U. Klaas (✉)

Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany
e-mail: klaas@mpia.de

K. Okumura

Commissariat à l’Energie Atomique, IRFU, Orme des Merisiers, Bât. 709, 91191 Gif/Yvette, France

M. Ferlet

Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire OX11 0QX, UK

T. Müller

Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße, 85748 Garching, Germany

M. Sanchez-Portal · B. Altieri

Herschel Science Centre, ESAC, Villanueva de la Cañada, 28691 Madrid, Spain

D. Doyle

ESA Scientific Projects Department, ESTEC/SRE-P, Keplerlaan 1,
2201 AZ Noordwijk, The Netherlands

G. L. Pilbratt

ESA Research and Scientific Support Department, ESTEC/SRE-SA, Keplerlaan 1,
2201 AZ Noordwijk, The Netherlands