

Reionization and the Cosmic Dawn with the Square Kilometre Array

Garrelt Mellema · Léon V. E. Koopmans · Filipe A. Abdalla · Gianni Bernardi · Benedetta Ciardi · Soobash Daiboo · A. G. de Bruyn · Kanan K. Datta · Heino Falcke · Andrea Ferrara · Ilian T. Iliev · Fabio Iocco · Vibor Jelić · Hannes Jensen · Ronniy Joseph · Panos Labrourpoulos · Avery Meiksin · Andrei Mesinger · André R. Offringa · V. N. Pandey · Jonathan R. Pritchard · Mario G. Santos · Dominik J. Schwarz · Benoit Semelin · Harish Vedantham · Sarod Yatawatta · Saleem Zaroubi

Received: 6 February 2013 / Accepted: 15 March 2013 / Published online: 27 April 2013
© Springer Science+Business Media Dordrecht 2013

Abstract The Square Kilometre Array (SKA) will have a low frequency component (SKA-low) which has as one of its main science goals the study of the redshifted 21 cm line from the earliest phases of star and galaxy formation in the Universe. This

G. Mellema (✉) · K. K. Datta · H. Jensen
Department of Astronomy & Oskar Klein Center,
Stockholm University, Stockholm, Sweden
e-mail: garrelt.mellema@astro.su.se

L. V. E. Koopmans · S. Daiboo · A. G. de Bruyn ·
R. Joseph · A. R. Offringa · H. Vedantham · S. Zaroubi
Kapteyn Astronomical Institute, University of Groningen,
Groningen, The Netherlands

F. A. Abdalla
Department of Physics and Astronomy,
University College London, London, UK

G. Bernardi
Center for Astrophysics, Harvard University, Cambridge, USA

B. Ciardi
Max Plank Institute for Astrophysics, Garching, Germany

A. G. de Bruyn · V. Jelić · P. Labrourpoulos · V. N. Pandey · S. Yatawatta
ASTRON, Dwingeloo, The Netherlands

H. Falcke
Department of Astronomy, Radboud University, Nijmegen, Netherlands

A. Ferrara · A. Mesinger
Scuola Normale Superiore, Pisa, Italy