

Herschel SPIRE FTS relative spectral response calibration

Trevor Fulton · Rosalind Hopwood · Jean-Paul Baluteau · Dominique Benielli · Peter Imhof · Tanya Lim · Nanyao Lu · Nicola Marchili · David Naylor · Edward Polehampton · Bruce Swinyard · Ivan Valtchanov

Received: 5 July 2013 / Accepted: 7 November 2013
© Springer Science+Business Media Dordrecht 2014

Abstract *Herschel/SPIRE* Fourier transform spectrometer (FTS) observations contain emission from both the *Herschel* Telescope and the SPIRE Instrument itself, both of which are typically orders of magnitude greater than the emission from the astronomical source, and must be removed in order to recover the source spectrum.

T. Fulton (✉) · P. Imhof
Bluesky Spectroscopy, Lethbridge, AB, T1J0N9, Canada
e-mail: trevor.fulton@blueskyinc.ca

T. Fulton · P. Imhof · D. Naylor · E. Polehampton
Institute for Space Imaging Science, Department of Physics & Astronomy, University of Lethbridge,
Lethbridge, AB, T1K3M4, Canada

R. Hopwood
Physics Department, Imperial College London, South Kensington Campus, London SW7 2AZ, UK

J.-P. Baluteau · D. Benielli
Aix Marseille Université, CNRS, LAM (Laboratoire d’Astrophysique de Marseille) UMR 7326,
13388 Marseille, France

T. Lim · E. Polehampton · B. Swinyard
RAL Space, Rutherford Appleton Laboratory, Chilton, Oxfordshire OX11 0QX, UK

N. Lu
NHSC/IPAC, 100-22 Caltech, Pasadena, CA 91125, USA

N. Marchili
Universitá di Padova, 35131 Padova, Italy

B. Swinyard
Department of Physics & Astronomy, University College London, Gower St, London WC1E 6BT, UK

I. Valtchanov
Herschel Science Centre, ESAC, 28691, P.O. Box 78, Villanueva de la Cañada, Madrid, Spain